




11/3/23

Outline and Objective

Anatomy and Physiology of the Vestibular System

- › Understand site of lesion and mode of presentation
- › Define Peripheral and Central Vestibular Disorders
- › Describe the Vestibulo-ocular Reflex


Aging of the Vestibular System



Part I Anatomy, Physiology and clinical implications


11/3/23

Peripheral Vestibular Anatomy and Physiology




11/3/23

Peripheral Vestibular Anatomy and Physiology




11/3/23




11/3/23

Peripheral Vestibular Anatomy and Physiology




11/3/23

Otolithic Membranes




11/3/23



11/3/23




11/3/23




11/3/23

Peripheral Vestibular Anatomy and Physiology



11/3/23

Courtesy of Michael
Teixido



org >

11/3/23



Labyrinth placed in 3D network with MRI based skull anatomy;
Amira 6.5 (Mercury Computer Systems, Chelmsford, MA, USA)



Labyrinth cloned / positioned relative to skull




org >

11/3/23



11/3/23

Peripheral Vestibular Anatomy and Physiology




(A) Adaptation is explained in the gating spring model by adjustment of the insertion point of tips links. Movement of the insertion point up or down the shank of the stereocilium, perhaps driven by a Ca^{2+} -dependent protein motor, can continually adjust the resting tension of the tip link. (After Hudspeth and Gillespie, 1994.)

[MUSChealth.org](https://www.muschealth.org)


11/3/23

Peripheral Vestibular Anatomy and Physiology

Kinocilium oriented toward Striola



Kinocilium oriented away from striola



MUSC Health
Medical University of South Carolina

Changing What's Possible | [MUSChealth.org](https://www.muschealth.org)

11/3/23




Figure 14.2

The morphological polarization of vestibular hair cells and the polarization maps of the vestibular organs. (A) A cross section of hair cells shows that the kinocilia of a group of hair cells are all located on the same side of the hair cell. The arrow indicates the direction of deflection that depolarizes the hair cell. (B) View looking down on the hair bundles. (C) In the ampulla located at the base of each semicircular canal, the hair bundles are oriented in the same direction. In the **sacculus** and **utricle**, the striola divides the hair cells into populations with opposing hair bundle polarities.

11/3/23




Figure 14.5

Forces acting on the head and the resulting displacement of the **otolithic membrane** of the utricle. For each of the positions and accelerations due to translational movements, some set of hair cells will be maximally excited, whereas another set will be maximally inhibited. Note that head tilts produce displacements similar to certain accelerations.




Figure 14.4

Morphological polarization of hair cells in the utricle and saccule. (A) Cross section of the utricle showing hair bundles projecting into the gelatinous layer when the head is level. (B) Cross section of the utricle showing the orientation of the kinocilia. (C) Orientation of the utricle and saccule in the head; arrows show orientation of the kinocilia, as in Figure 14.2. The saccule on either side are oriented more or less vertically, and the utricle more or less horizontally. The striola is a structural landmark consisting of small stereocilia arranged in a narrow trench that divides each **otolith** organ. In the utricle macula, the kinocilia are directed toward the striola. In the saccular macula, the kinocilia point away from the striola. Note that, given the **utricle** and **sacculus** on both sides of the body, there is a continuous representation of all directions of body movement.

11/3/23

Remember: Both vestibular systems are working when we are not moving (push/pull mechanism)



11/3/23

Remember: Both vestibular systems are working when we are not moving




collab
clips



11/3/23

Semicircular canals and Balance

Flourens 1842 Ewald 1892



11/3/23



Ewald's laws




EWALD'S THREE LAWS

1. A stimulation of the semicircular canal causes a movement of the eyes in the plane of the stimulated canal
: **tonic movements**(slow phase of nystagmus and segmental and axial tilts) are in the **direction of the endolymph movement**
2. In the **horizontal** semicircular canals, an **ampullopetal** endolymph movement causes a **greater stimulation** than an ampulofugal one. **Ampullopetal flow is utriculopetal**
3. In the **vertical** semicircular canals, the reverse is true. **Ampullopetal flow is utriculofugal**

Robert Bárány XXth Century



Head Impulse Test (bHIT)**



**Petersen JA, Straumann D and Weber K, Clinical Diagnosis of bilateral vestibular loss: three simple bedside tests, Ther Adv Neurol Disord, 2013,6: 41-45

11/3/23

Head Impulse Test (bHIT)**



MUSC Health
Medical University of South Carolina

Changing What's Possible | MUSChealth.org



VOR SUPPRESSION by visual fixation

**Brandt T and Strupp M, General
Vestibular Testing, Clin Neurophys,
2005, 116: 406-426**


MUSC Health
Medical University of South Carolina

Changing What's Possible | MUSChealth.org



11/3/23

Velocity Storage Asymmetry



11/3/23

Dix, M.R.; Hallpike, C.S.: The pathology, symptomatology and diagnosis of certain common disorders of the vestibular system. Proc. R. Soc. Med. 45: 341–354 (1952).





FIG. 7.

By means of apparatus shown in Fig. 7 it is possible to move the patient *en masse* into the critical position,

11/3/23




	Excitatory Pathway	Inhibitory Pathway
Superior SCC	- Bilateral Superior Recti - Contralateral Inferior oblique	- Bilateral Inferior Recti - Contralateral Superior Oblique
Posterior SCC	- Ipsilateral superior oblique - Bilateral Inferior Recti	- Ipsilateral inferior oblique - Bilateral Superior Recti
Lateral SCC	- Contralateral lateral Rectus - Ipsilateral Medial Rectus	- Contralateral Medial Rectus - Ipsilateral Lateral Rectus

USCHealth.org >

11/3/23

Posterior Canal BPPV




Superior oblique muscle: depressor when eye is adducted and intorter when abducted
Left Posterior BPPV

NEUTRAL GAZE: TYPICAL UPBEAT GEOTROPIC NYSTAGMUS WITH VERTICAL AND ROTARY NYSTAGMUS
LEFT GAZE: PURE GEOTROPIC NYSTAGMUS ROTARY COMPONENT ONLY
RIGHT GAZE: PURE UPBEAT NYSTAGMUS VERTICAL COMPONENT ONLY

MUSC Health
Medical University of South Carolina

Changing What's Possible | MUSCHealth.org >


11/3/23



Arch
Otolaryngol
1969: 90, 765


11/3/23

Buoyancy Cupulopathy



11/3/23

Superior canal Bppv



Excitation of right and left superior recti and inhibition of right and left inferior recti
 Excitation of left contralateral inferior oblique and inhibition of left contralateral superior oblique:
 ELEVATES THE LEFT EYE WHEN IT IS ADDUCTED

MUSC Health
 Medical University of South Carolina

Changing What's Possible | MUSChealth.org >

11/3/23

Lateral Canal BPPV



MUSC Health
 Medical University of South Carolina

Changing What's Possible | MUSChealth.org >

11/3/23

Define the symptoms

Lightheadedness/Presyncope	- Sensation of impending loss of consciousness - Multiple sensory disturbances, orthostatic hypotension, cardiac arrhythmias and panic attacks
Disequilibrium	- Imbalance or unsteadiness experience while standing or walking. - Caused by diminished vision, loss of vestibular function, defects in proprioception and motor dysfunction from the CNS and PNS. - Also related to visual vertigo, presyncopal faintness and somatoform phobic postural vertigo
Oscillopsia	- Subjective illusion of visual motion. - Happens ONLY when eyes are open: - Acquired nystagmus can cause retinal slip. - Bilateral vestibular loss
Vertigo	Illusion of movements of oneself or the environment: rotationally but not necessarily

Huh YE, Kim JS, Bedside Evaluation of dizzy patients, J Clin Neurol, 2013,9: 203-213

MUSC Health
Medical University of South Carolina

Changing What's Possible | MUSChealth.org



11/3/23

Oscillopsia



MUSC Health
Medical University of South Carolina

Changing What's Possible | MUSChealth.org



11/3/23



Journal of Vestibular Research 19 (2009) 1–13
 DOI:10.3233/JVES-2009-0343
 IOS Press

1

Classification of vestibular symptoms: Towards an international classification of vestibular disorders

First consensus document of the Committee for the Classification of Vestibular Disorders of the Bárány Society

Alexandre Bisdorff^{a,*}, Michael Von Brevern^b, Thomas Lempert^c and David E. Newman-Toker^d

^aDepartment of Neurology, Centre Hospitalier Emile Mayrisch, L-4005 Esch-sur-Alzette, Luxembourg

^bVestibular Research Group Berlin, Department of Neurology, Park-Klinik Weissensee, Berlin, Germany

^cVestibular Research Group Berlin, Department of Neurology, Schlosspark-Klinik, Berlin, Germany


^dDepartment of Neurology, The Johns Hopkins University School of Medicine, Baltimore, MD 21287, USA

On behalf of the Committee for the Classification of Vestibular Disorders of the Bárány Society: Pierre Bertholon, Alexandre Bisdorff, Adolfo Bronstein, Herman Kingma, Thomas Lempert, Jose Antonio Lopez, Escamez, Måns Magnusson, Lloyd B. Minor, David E. Newman-Toker, Nicolás Pérez, Philippe Perrin, Mamoru Suzuki, Michael von Brevern, John Waterston and Toshiaki Yagi






Changing What's Possible | MUSChealth.org

Appendix 2. Symptom Coding Algorithm




Changing What's Possible | MUSChealth.org






11/3/23

Peripheral Vestibular Anatomy and Physiology



11/3/23

Physiology of Endolymph



Leveque M, Seidermann L, Ulmer E and Chays A, [Vestibular Physiology: anatomic, cellular immunohistochemical and molecular substrates], EMC, 2010.

MUSC Health
University of South Carolina

Changing What's Possible | MUSChealth.org

11/3/23

Tumarkin crisis

Tumarkin A. The otolithic catastrophe. A new syndrome. Br Med J 1936;1:175-7

- Utricle affection: accessions of rigidity or flexor spasm
 - "Mr X was standing at his desk talking to a client when suddenly he slumped to the floor. He had no vertigo, no loss of consciousness and no malaise. It The thing came like a bolt from the blue but he was immediately able to assure onlookers he was all right and almost immediately got up and carried on."
- Saccular affection: Fall to the side
 - "Mr Y was sent to me on account of giddiness.[...] On one occasion he was standing in his van receiving packages when suddenly he felt as if the wheel had collapsed under him."

	Canal	Vestibule
Onset	Ingravescent	Instantaneous
Duration	May be minutes up to days	A minute or less
Nausea, pallor, sweating, etc.	Yes	No
Giddiness	Yes	No
Deafness	Variable in type: may be little or none	Definite reduction of bone conduction




FIG. 1a AND 1b. FIG. 2. FIG. 3.
In the upright position (Fig. 1a) b is pressing on a. In the upside-down position (Fig. 1b) it will drag, and intermediate positions will produce intermediate effects. The right and left lapilli are synergic.

Sudden shift of utricle macula, sudden changes in EL pressure, sudden electrolyte changes from rupture of membrane




FIG. 4. FIG. 5.
The sensitive area (sagitta) lies on the side instead of the floor of its retaining sac; furthermore the right and left sagittae are roughly at right angles to each other, so that unlike the lapilli they are opponents and not synergists.

What's Possible | MUSChealth.org

11/3/23

Tumarkin Crisis Post-Epley



MUSC Health
Medical University of South Carolina

Changing What's Possible | MUSChealth.org



11/3/23

Are we gravity bound?

Caloric experiments in microgravity

- › Convection currents not present
- › Thermal effect

Antigravity vector “built-in” in otolith/graviceptive pathways.

Pierrot-Deseilligny C, Effect of Gravity on Vertical Eye
Position, Ann NY Acad Sci, 2009, 1164: 155-165

MUSC Health
Medical University of South Carolina

Changing What's Possible | MUSChealth.org




11/3/23

Parabolic Flight “The Vomit Comet”



11/3/23

Are we gravity bound?



11/3/23


When astronauts return home they experience some after effects from this otolith organ disruption:

- › Difficulty walking around corner
- › Sensation of linear movement or freefall when tilting one's head

Vestibular neuronitis can cause similar symptoms

11/3/23

Vestibular System Pleiotropic Mediator



11/3/23

Central Vestibular Anatomy and Physiology

Cerebellum: major role in motor coordination


Archeocerebellum or Vestibulocerebellum: flocculus, nodulus, vermis, fastigial nucleus

- › Coordinate movements of head and eyes
- › Motor learning, adaptation and compensation

Paleocerebellum or Spinocerebellum

11/3/23

FUNCTIONAL ORGANIZATION OF CEREBELLUM




Physiological vertigo	Pathological Vertigo	Vestibular Function	Vertigo Syndrome
Vestibular	Peripheral labyrinthine lesion Peripheral eighth nerve lesion Central vestibular lesion	Spatial orientation Motion perception	Vertigo
Optokinetic		Vestibulo-Ocular Reflex	Nystagmus
Somatokinetic		Posture	Ataxia
		Vegetative effects	Nausea

Brandt T, Daroff RB. The
and pathological vertigo
1980;7:194–203.

MUSC Health
Medical University of South Carolina

Changing What's Possible | MUSChealth.org

Figure 14-15a Motor and Sensory Regions of the Cerebral Cortex



© 2012 Pearson Education, Inc.





Figure 13.14

The human **auditory cortex**. (A) Diagram showing the brain in left lateral view, including the depths of the lateral sulcus, where part of the **auditory cortex** occupying the superior temporal gyrus normally lies hidden. The primary **auditory cortex** (A1) is shown in blue; the surrounding belt areas of the **auditory cortex** are in red. (B) The primary **auditory cortex** has a tonotopic organization, as shown in this blowup diagram of a segment of A1.

Homunculus of Primary Somatosensory Cortex in Blue

Note that each hemisphere receives info from the opposite side of the body



MUSC Health
Medical University of South Carolina

Changing What's Possible | MUSChealth.org

Area 7 – Visuo-Motor Coordination
Point of convergence between vision and proprioception to determine where moving objects are in relation to the body

Area 39 – Angular gyrus, considered by some to be part of **Wernicke's area**
It is also involved in a number of processes related to language, number processing and **spatial cognition**, memory retrieval, **attention**, and **theory of mind**.

Area 40 – Supramarginal gyrus, involved with language processing. Lesions in it can result in receptive aphasia

Vestibular cortices and spatial cognition. Vestibular cortices involved in spatial cognition, illustrated in a soccer player: HD cell, head direction cell; HPC, hippocampus; MST, medial superior temporal area; ParaHPC, parahippocampal cortex; Parietal Ant, parietal anterior cortex; PIVC, parieto-insular vestibular cortex; RSP, retrosplenial cortex; VIP, ventral intraparietal area; 7a, 39, 40 Brodmann area.

MUSC Health
Medical University of South Carolina

Changing What's Possible | MUSCHealth.org

11/3/23

Central Vestibular Anatomy and Physiology

Vestibulothalamic and vestibulocortical pathways:

- › **sensation and perception of head movement**

Vestibulospinal pathways:

- › **motor commands to the muscles of the neck, upper torso and lower limbs to maintain balance**
- › **Cortical efferent and cerebellar efferents modulate vestibulospinal output**
- › **Related to drop attacks**

Vestibulocular pathways

- › **responsible for the VOR**
- › Responsible for gaze, saccades, smooth pursuit and optokinetic nystagmus

11/3/23

Two questions come to mind

Q1 Can someone recover from damage to the end-organ?



Vestibular Neuronitis: Unilateral Vestibular Deafferentation

34 year old patient with no prior medical problems

6/6/2017 felt spinning when he would lay on his left side. This gets better when he lays flat.

6/8/2017 Acute right stye

6/10/2017 vertigo constant even when looking straight ahead



11/3/23

Alexander's Law: 1912

In Unilateral vestibular lesion


- › Spontaneous Nystagmus with fast phase toward healthy ear

Slow phase is greatest when gaze is directed toward healthy ear or nystagmus fast phase, attenuates at central gaze and weakest when gaze is toward ipsilesioned ear

It actually is a compensation strategy from CNS in the context of vestibular imbalance

11/3/23


VideoHIT




11/3/23

Vertigo starts subsiding and patient starts noting falling to the right side when he turns his head to the right or his ear to touch his shoulder

11/3/23




11/3/23




11/3/23

Patient's balance is off for the next ten days
No more spinning vertigo




11/3/23



Changing What's Possible | MUSCHealth.org


11/3/23

36 days s/p onset




Changing What's Possible | MUSCHealth.org

Vestibular Test Battery		Evaluation Timepoints					
		Week 1	Week 2	Week 3	Week 6	Week16	Month 8
Calorics			41% RUW	47% RUW	78% RUW	0% BW	59% RUW
Spontaneous Nystagmus			3 d/s LBN	None	None	None	None
SHA gain			Normal	Normal			Normal
SHA symmetry			CW weaker	Normal			Normal
SHA phase			Normal	Normal			Normal
vHIT gain	Right	0.77	0.23	0.29	0.33	0.60	0.82
	Left	0.95	0.59	0.63	0.8	0.59	0.83
vHIT overt saccades	Right	Present	Present	Present	Present	Present	Present
	Left	Absent	Absent	Absent	Absent	Absent	Present
vHIT covert saccades	Right	Present	Present	Present	Present	Absent	Absent
	Left	Absent	Absent	Absent	Absent	Absent	Absent




9/6/2017
Day 90




Name: Habib Risk
 Birthday: 008
 Test: Head Impulse
 Variant: Lateral
 Date: 2017-09-06 / 15:56:53
 Examiner:
 Calibration: Eye: default / Head: default


9/20/2019
3.5 Month



Name: Demo Demo
 Birthday: 1970-05-31
 Pans ID: 0101013333
 Test: Head Impulse
 Variant: Lateral
 Date: 2017-09-20 / 10:23:11
 Examiner:
 Calibration: / Head: default



Month 8



Bilateral Vestibular Deafferentation

Sequential:

- › Bechterew's Phenomenon

Concomitant

March 20, 1952

“Is there any man-made machine designed like the human apparatus-with so many alternate systems to accomplish its end?”



11/3/23

Stimulator


Vestibular electrode (PAN)

Vestibular electrode (LAN)


Vestibular electrode (SAN)

Intracochlear array

A



B



C



11/3/23

Audiology &
Neurotology

Review

Audiol Neurotol 2020;25:6–24
DOI: 10.1159/000502407Received: May 31, 2019
Accepted after revision: July 29, 2019
Published online: September 18, 2019**Electrical Vestibular Stimulation in Humans: A Narrative Review**Morgana Sluydts^a Ian Curthoys^b Robby Vanspauwen^a Blake Croll Papsin^c
Sharon Lynn Cushing^c Angel Ramos^d Angel Ramos de Miguel^d
Silvia Borkoski Barreiro^d Maurizio Barbara^e Manuel Manrique^f
Andrzej Zarowski^a^aEuropean Institute for Otorhinolaryngology, GZA Hospitals Antwerp, Wilrijk, Belgium; ^bVestibular Research Laboratory, University of Sydney, Sydney, NSW, Australia; ^cDepartment of Otolaryngology – Head and Neck Surgery, The Hospital for Sick Children, Toronto, ON, Canada; ^dHearing Loss Unit, Otorhinolaryngology, Head and Neck Department, Complejo Hospitalario Universitario Insular Materno Infantil, Las Palmas de Gran Canaria, Spain; ^eNESMOS Department, ENT Clinic, Sapienza University, Rome, Italy; ^fOtorhinolaryngology Department, Clínica Universidad de Navarra, Pamplona, SpainMUSC Health
Medical University of South Carolina

Changing What's Possible | MUSCHealth.org >

11/3/23

- Vestibular costimulation with a cochlear implant
- Human vestibular implant
 - Vestibular Pacemaker-Vestibular neurostimulator (Rubenstein University of Washington): 2 extracochlear electrodes and 3 SCC electrodes in perilymphatic space
 - Vestibulocochlear implant (Maastricht-Geneva): intracochlear electrode and 1-3 SCC electrodes
 - Multichannel Vestibular implant (Della Santina Johns Hopkins): built-in gyroscopes and accelerometers Fifty electrodes over three shanks: one shank for saccule and horizontal SCC, one shank for utricle and superior SCC and one separate shank for the posterior SCC
 - Otolith Implant (Ramos de Miguel University of Las Palmas) insertion into the vestibule

MUSC Health
Medical University of South Carolina

Changing What's Possible | MUSCHealth.org >

11/3/23

Q2 Can the peripheral vestibular end-organ affect higher cortical functions?



Changing What's Possible | MUSChealth.org



PART II The vestibular system-
cognition link

11/3/23

Aging of the Vestibular System Histology

Marked age-related degeneration in nearly every type of vestibular-related cell and neuron bundle

- › Starting after the age of 65 years
- › Saccule seems to be the most affected

Clinical Implications: Age-related declines in the ability to correctly detect head position and motion in space, to elicit vestibular spinal reflexes or to solve sensory conflicts

Tang PF, Woollacott M, Balance control in older adults. In: Bronstein AM, Brandt TW, Nitt J, eds. Clinical disorders of balance, posture and gait. Amsterdam: Elsevier; 2004:385-403



Changing What's Possible | MUSCHealth.org



11/3/23

Aging of the Vestibular System Clinical Correlation

Early studies did not show direct correlation between histology and abnormalities on paraclinical testing

More recent evidence comes with VEMPs and videoHIT showing high prevalence of abnormalities with age

1. Agrawal Y, Zuniga MG, Davalos-Bichara M et al, Decline in semicircular canal and otolith function with age, *Otol Neurotol*, 2012; 33: 832-839
2. Agrawal Y, Schubert MC, Migliaccio AA et al, Evaluation of quantitative head impulse testing using search coils versus video-oculography in older individuals, *Otol Neurotol*, 2014; 35:283-288

Central compensation occurs until a “tipping point” is reached



Changing What's Possible | MUSCHealth.org



11/3/23

Aging of the Vestibular System Epidemiology

Incidence of Dizziness increases from 22% (65-69) to over 40% for adults over 80-84 (18 M people by 2050)

Cutson TM, Falls in the elderly, Ann Fam Physician, 1994, 49:149-156

65% of adults over 60 experience dizziness and loss of balance on a daily basis

Hobeika CP, Equilibrium and balance in the elderly, Ear Nose Throat J, 1999, 78:558-562

Approximately 20% of imbalance and 40-50% of dizziness is due to a vestibular problem

University of Iowa Health Care. Comprehensive Management of vestibular disorders, Currents, 2002,3 (2)

11/3/23

Aging of the Vestibular System Epidemiology

TABLE 1.1
Most Common Causes of Dizziness in Primary Care Practice¹

Category	Percentage of Patients (%)	Examples
Peripheral vestibular disease	20–50	Benign paroxysmal positional vertigo, labyrinthitis, vestibular neuritis
Cardiovascular disease	10–30	Arrhythmia, congestive heart failure, vasovagal conditions (e.g., carotid sinus hypersensitivity)
Systemic infection	10–20	Systemic viral and bacterial infection
Psychiatric conditions	5–15	Depression, anxiety, hyperventilation
Metabolic disturbances	5–10	Hypoglycemia, hyperglycemia, electrolyte disturbances, thyrotoxicosis, anemia
Medications	5–10	Antihypertensives, psychotropic medications

11/3/23

Aging of the Vestibular System Epidemiology

Lifetime prevalence of vestibular vertigo 7.8%

One year prevalence of vertigo 4.9%

- › 7.2% in 60-69 year olds
- › 8.8% over the age of 80

Incidence of vertigo 1.5%

Neuhauser NK, von Brevern M, Radtke A et al. Epidemiology of vestibular vertigo: a neurologic survey of the general population. Neurology, 2005; 65:898-904



Changing What's Possible | MUSChealth.org



11/3/23

Aging of the Vestibular System Epidemiology

35% of US adults over 40 have evidence of balance dysfunction on postural exam

85% of individuals over 80 have evidence of balance dysfunction

Agrawal Y, Carey JP, Della Santina CC, Schubert MC, Minor LB, Disorders of balance and vestibular function in US adults: data from the national health and nutrition examination survey, 2001-2004, Arch Intern Med, 2009;169:938-944

Dizziness is a major cause of absenteeism, partial or full disability and an increase in healthcare utilization



Changing What's Possible | MUSChealth.org



11/3/23

Diagnostic criteria for presbyvestibulopathy (PVP)

Each of the criteria A through D have to be fulfilled

A. Chronic vestibular syndrome (at least 3 months duration) with at least 2 of the following symptoms:¹

1. Postural imbalance or unsteadiness
2. Gait disturbance
3. Chronic dizziness
4. Recurrent falls

B. Mild² bilateral peripheral vestibular hypofunction documented by at least 1 of the following:

1. VOR gain measured by video-HIT between 0.6 and 0.8³ bilaterally
2. VOR gain between 0.1 and 0.3 upon sinusoidal stimulation on a rotatory chair (0.1 Hz, Vmax = 50-60°/sec)⁴
3. Reduced caloric response (sum of bithermal maximum peak SPV on each side between 6 and 25°/sec)⁵

C. Age ≥ 60 years⁶

D. Not better accounted for by another disease or disorder⁷

11/3/23

Presbyvestibulopathy




FIG. 1.1 Typology of age-related dizziness, imbalance, and vestibular loss. VOR, vestibulo-ocular reflex; VSR, vestibulospinal reflex.

11/3/23



11/3/23

Vestibular System and Cognition

Older studies:

- › Memory impairment and Perilymphatic fistulae
- › Concentration difficulties and gentamicin ototoxicity

Bilateral vestibular nerve section (NF2) and reduction in spatial memory and spatial navigation as well as reduced hippocampal volumes

Brandt T, Schautzer F, Hamilton DA, Brüning R, Markowitsch HJ, Kalla R, Strupp M, Vestibular loss causes hippocampal atrophy and impaired spatial memory in humans, Brain, 2005, 128:2732-2741

11/3/23

Van der Zaag HJ, van Leeuwen RB, Dizziness causes absence from work, Acta Neurol Belg, 2015, 115: 345-349

- › 400 patients
- › 70% absenteeism
- › 12% completely disabled
- › 51% worked less because of the dizziness



11/3/23

Vestibular System and Cognition

Box 1. Summary of 3 US population-based studies reporting a significant association between reduced vestibular function and poorer cognitive performance.

Study population	Year	N	Finding
Baltimore Longitudinal Study of Aging (BLSA)	2013-2014	183	Reduced saccular function associated with significantly poorer scores on neurocognitive tests of visuospatial ability in cross-sectional analyses; no significant relationship between saccular function and tests of language, executive function, and verbal memory
National Health and Nutrition Examination Survey (NHANES)	1999-2002	1303	Vestibular impairment (based on modified Romberg test) associated with significantly poorer Digit Symbol Substitution Score in cross-sectional analyses
National Health Interview Survey (NHIS)	2008	20,950	Vestibular vertigo (based on questionnaire responses) significantly associated with 4-fold increased odds of "difficulty remembering and confusion" in cross-sectional analyses

Bigelow, R. T., Semenov, Y. R., Du Lac, S., Hoffman, H. J., & Agrawal, Y. (2015). Vestibular vertigo and comorbid cognitive and psychiatric impairment: The 2008 national health interview survey. *Journal of Neurology, Neurosurgery & Psychiatry*, 87(4), 310–319.

Bigelow, R. T., Semenov, Y. R., Trevino, C., Ferrucci, L., Resnick, S. M., Simonsick, E. M., ... Agrawal, Y. (2015). Association between visuospatial ability and vestibular function in the Baltimore longitudinal study of aging. *Journal of the American Geriatrics Society*, 63(9), 1837–1844.

Semenov, Y. R., Bigelow, R. T., Xue, Q., Lac, S. D., & Agrawal, Y. (2015). Association between vestibular and cognitive function in US adults: Data from the national health and nutrition examination survey. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, 71(2), 243–250.



11/3/23

Vestibular System and Cognition

Hippocampal atrophy in Meniere's disease

1. Kremmyda O, Hüffner K, Flanagan VL, Hamilton DA, Linn J, Strupp M, Brandt T, *Beyond dizziness: Virtual navigation, spatial anxiety and hippocampal volume in bilateral vestibulopathy, Frontiers in Human Neuroscience*, 10, 139
2. Popp P, Wulff M, Finke K, Rühl M, Brandt T and Dieterich M, *Cognitive deficits in patients with chronic vestibular failure, J Neurol*, 2017, 264:554-563

20% of patients presenting with vertigo have reproducible and significant cognitive error in arithmetic abilities, short term memory and performance in specific visual environments

1. Risey J and Briner W, *Dyscalculia in Patients with Vertigo, J Vest Res*, 1990, 1:31-37
2. Hanes DA and McCollum G, *Cognitive-vestibular interactions: a review of patient difficulties and possible mechanisms, J Vest Res*, 2006, 16:75-91



Changing What's Possible | MUSChealth.org



11/3/23

Vestibular System and Cognition

Alzheimer's disease patients: twice as likely to have vestibular impairment than healthy controls

AD patients with vestibular impairment have disproportionate reduction in spatial cognition compared to AD without vestibular loss

- Agrawal Y, Smith PF and Rosenberg PB, Vestibular impairment, cognitive decline and Alzheimer's disease: balancing the evidence, Aging and Mental Health.*
Doi:10.1080/13607863.2019.1566813



Changing What's Possible | MUSChealth.org



11/3/23





Figure 1. Conceptual model of impact of aging on vestibular function (notably saccular function), which contributes to neurodegeneration of neural circuits involved in vestibular processing and deterioration specifically in spatial cognitive ability.

11/3/23

Peripheral vestibular system provides major inputs to cholinergic neurons in medial temporal region, specifically degraded in AD

- Vestibular impairment may participate in pathogenesis of AD
- Vestibular Impairment may cause a spatial phenotype of AD where there is higher risk of spatial disorientation, wandering and increase fall risk



11/3/23

Wide dissemination of vestibular signals: maybe an evolutionary process to create a neural network with sparse coding

Sparse coding:

- › Low energy cost
- › Large storage capacity
- › Rapid learning ability,
- › Tolerance to degradation



11/3/23

Higher vestibular function

We are taught to clinically think in terms of peripheral versus central based on anatomical lesions

There is a lot of overlap in clinical presentations

Brandt T, Strupp M and Dieterich M, Toward a concept of disorders of "higher vestibular function", Front Integ Neuroscience, 2014,8 doi:10.3389/fnint.2014.00047



11/3/25




FIGURE 1 | This figure proposes a collection of clinical syndromes which may be called central vestibular disorders or disorders of higher vestibular function. They are depicted in alphabetical order and topographically grouped for cerebral cortex, thalamus, brainstem and cerebellum. The topographic assignment remains uncertain for some conditions. Note also that similar disorders occur with lesions at different sites—brainstem or cortex (room tilt illusion) or brainstem and cerebellum (ocular tilt reaction (OTR))—within the central vestibular neuronal circuitry. Please note that this list does not include all central vestibular syndromes. VC = vestibular cortex; VT = vestibular thalamus; VN = vestibular nucleus; VL = vestibular labyrinth.

11/3/23

Thank you for listening

Feel free to email me questions at rizkh@Musc.edu

You can also reach me on my cellphone 2154590435

SHARE YOUR VESTIBULAR EXPERIENCE

Add your information to VEDA's Patient Registry to help vestibular researchers and advocates

YOUR STORY IS MORE POWERFUL THAN YOU THINK!

HOW YOU CAN HELP

- 1 Go to vestibular.org/registry
- 2 Create an account
- 3 Fill out the surveys
- 4 Your information is off to work!

Questions?
Call 1 (800) 837-6428 or
Email info@vestibular.org

VEDA
VESTIBULAR.ORG

Charleston Dizzy Group

<http://www.facebook.com/groups/542582642742649>

MUSC Health
Medical University of South Carolina

Changing What's Possible | MUSChealth.org